

Effect of Trace Elements from Consumption of Crops Cultivated on Mine Domes (Spoils): A Case Study of Jos Plateau and Environs, Central Nigeria

Raymond Revelation Dakar¹, Umaru Umaru Nahabu¹, Adamu Yunusa Ugya², Tijjani Sabiu Imam^{2,*}

¹Federal College of Animal Health and Production Technology, National Veterinary Research Institute, Vom, Jos, Nigeria

²Biological Sciences Department, Bayero University, Kano, Nigeria

Email address:

tsimam.bio@buk.edu.ng (T. S. Imam)

*Corresponding author

To cite this article:

Raymond Revelation Dakar, Umaru Umaru Nahabu, Adamu Yunusa Ugya, Tijjani Sabiu Imam. Effect of Trace Elements from Consumption of Crops Cultivated on Mine Domes (Spoils): A Case Study of Jos Plateau and Environs, Central Nigeria. *International Journal of Pharmacy and Chemistry*. Vol. 3, No. 5, 2017, pp. 67-71. doi: 10.11648/j.ijpc.20170305.12

Received: May 30, 2017; **Accepted:** August 26, 2017; **Published:** October 26, 2017


Abstract: The study was conducted on the Jos Plateau metropoly and its environs in Jos North, parts of Jos South and Barkin-Ladi Local Government Areas of Plateau State, Central Nigeria. The studied area covered south eastern part of Naraguta sheet 168 NE, extends down North eastern part of Kurra sheet 189NE bounded by Longitude 8°50'E to 9°00'E and Latitude 9° 26'N to 9°43'N respectively. This area sampled covers a land mass of about 586.25 square kilometer. This study begins from May, 2009 to March, 2016 at the National Veterinary Research Institute (NVRI), Vom. The aim was to investigate the effect of trace elements from consumption of crops cultivated on mine domes (spoils). The results presented in this paper refer to Tomato and Irish-potato, which are the most popular crops planted in that area in 2009-2016 respectively. Energy Dispersive X-Ray Fluorescence spectrometry(ED-XRFS) method was employed for the analysis of Cu, Fe, Zn, Ni, Pb, Cr and Mn. This study concluded that all those concentrations are well below threshold limit values accepted in Nigeria. No significant differences in Cu, Pb and Cr concentrations between the tested Tomato and Irish-potato were found. Zinc concentration in Potato was significantly lower than in Irish potato. Generally, the study concluded that plant samples were found to have major, minor and mostly trace elements' concentrations below the safe values. However, Pb that has high concentration above the safe value (threshold limit) in Potato is advised not to be cultivated around mine domes, because of its health adverse effects on the inhabitants.

Keywords: Mine, Domes, Concentrates, Trace Elements, Tomatoes, Irish Potatoes

1. Introduction

Trace elements play a significant role in many metabolic processes in plant organisms, but at excessive concentrations, particularly in soluble forms and easily available to plants, can cause phytotoxic symptoms. Plant response to high contamination of soil and air with heavy metals is extremely variable and depends on many environmental factors [1] [2]. Soluble and colloidal forms of heavy metals suspended in a soil solution are the most important for plant uptake, and the most important factors

influencing their availability and their total concentration and soil reaction [3]. The accumulation of heavy metals and metalloids in agricultural soil is of increasing concern nowadays. Potentially harmful metal in soil may come from the bedrock itself and anthropogenic sources like solid or liquid waste deposits, agricultural inputs and fallout of industrial and urban emissions [4]. Excessive accumulation in agricultural soils results in soil contamination and has consequences for food quality and safety. Food is the major



Source: National Centre for Remote Sensing Plateau State, Nigeria.

Figure 1. Geological map showing Locations of sampling sites for Tomato and Irish potato.

Table 2. Mean concentrations of each element analyzed from Irish potato sample in (mg/kg).

| S/No | Elements | Concentrations in (mg/kg) of irish potato samples | | | | | | | mean | Range Min Max |
|------|----------|---|-------|-------|-------|-------|-------|-------|------------|---------------|
| 1 | Cu | 96.1 | 103.9 | 80.7 | 86.3 | 73.5 | 71.1 | 82.2 | 71.1-103.9 | |
| 2 | Fe | 148.3 | 29.4 | 278.4 | 183.2 | 504.9 | 211.9 | 226.0 | 29.4-504.9 | |
| 3 | Zn | 21.7 | 20.1 | 16.1 | 17.7 | 14.5 | 72.3 | 27.0 | 14.5-72.3 | |
| 4 | Cr | 0.4 | 1.4 | 13.7 | 12.3 | 15.7 | 2.7 | 52.2 | 0.4-15.7 | |
| 5 | Mn | 17.0 | 32.5 | 33.3 | 24.8 | 37.2 | 17.0 | 26.9 | 17.0-37.2 | |
| 6 | Ni | 18.1 | 15.9 | 9.5 | 11.0 | 12.6 | 12.6 | 13.3 | 9.5-18.1 | |
| 7 | Pb | ND | ND | 1766 | 1377 | 1740 | ND | 1627 | ND-1766 | |

Table 3. Ranges of elements and their crustal abundance in plants.

| Names of elements | Ranges of plant concentration in (mg/kg) | | Safe | Crustal abundance (Mg/kg) |
|-------------------|--|------------|------|---------------------------|
| | Min. Range | Max. Range | | |
| Cu | 7-30 | 84-17000 | 1000 | 8.7 |
| Zn | 21-70 | 101-49000 | 2500 | 11 |
| Mn | 31-100 | 32-9870 | - | 170 |
| Fe | 21-70 | - | - | 10,000 |
| Cr | 30 | 480-1300 | 200 | 19 |
| Ni | 56 | 106-544000 | 300 | 13 |
| Pb | - | 1200 | - | - |

Note: Where (ND) = Not Detectable. Sources: [11] [12].




Figure 2. Graph showing the concentration of trace elements in tomato samples in mg/kg.




Figure 3. Graph showing the concentration of trace elements in Irish potato samples in mg/kg.

All the plants (Tomato and Irish potato) in the study area are summarized in (Tables 1 and 2) and in the graphs (2 and 3), display a very similar distribution pattern of trace elements. These similarities suggest that, the geologic processes associated with the release of these elements from the host rock and soil is the same. The distribution of trace element in tomato samples as shown in Table 3, when compared with the concentration of plants, shows that Cu, Zn, Mn, Fe, Cr, Ni except Pb that was not detectable (ND) possesses values below the safe or permissible concentrations. Similarly, Table 2 shows that Cu, Zn, Mn, Fe, Cr and Ni except Pb exhibit the permissible concentration [7] [12].

Copper (Cu); going by the safe concentration. 1000mg/kg in (Table 3), the arithmetic mean values of copper from the analysis in both Tomato and Irish potato is 76.2 mg/kg and 82.2 mg/kg respectively from (Table 1 and 2), shows that the concentration of copper within the study area is at a non objectionable rate in plant [2] [7].

The arithmetic mean values of Fe from the analysis of both Tomato and Irish potato as shown in Tables 1 and 2 having 235.3 mg/kg and 211.9mg/kg values respectively. These, when correlated with its crustal abundance in Table 3, 10,000mg/kg. This showed that, the study area has high Fe concentration [2].

Cr and Ni concentrations are permissible both in Tomato and Irish Potato (Table 1 and 2). Their Arithmetic mean values in tomato and irish potato ranges from 20.3 – 52.2 Cr and 7.2 – 13.2 Ni. These values are safe in both plants, except where it goes beyond the safe value in Table 3. This could cause cancer in man and would be so toxic to Plants when it exceeds the safe values.

The concentration of Pb 1,627 mg/kg in Irish potato exceeded the recommended value of 1,200mg/kg (Table 3). This indicates that, the study area are cultivating and consuming objectionable concentration of Pb in the Irish potato. Even though there was no concentration of this metal was detected in tomato crop. However, the arithmetic mean

values of Mn, in tomato and Irish potato, 67.5mg/kg and 26.9mg/kg respectively as compared to the standard in Table 3, shows that, the study area has the minimum concentration of this trace element [13].

In general the plant samples were found to have major, minor and mostly trace elements' concentrations below the safe value as shown in Table 3, however, Pb that has high concentration above the safe value (threshold limit) in the Irish potato is advised not to be cultivated, because of its adverse health effects on the inhabitants [14].

4. Conclusions

In general the plant samples were found to have major, minor and mostly trace elements' concentrations below the safe value. It is note worthy that irish potato accumulated Pb above the threshold limit which makes unsuitable for consumption.

The anomalous concentration of trace elements in the analyzed plant crops samples gives evidence of extensive mineralization and also the anomalous concentration of some trace elements like Pb, Fe, Cr, Ni, Mn, etc. may also suggested that there is every tendency for the water to be affected by these elements due to remobilization into the water phase. The presence of gorges, abandoned mine, dams, mine pits, constitute a huge threat to the inhabitants of the area.

The research indicates also that, the prevalent high rate of cancer in the northern (Upper) part of Plateau State is as a result of high rate of cultivation and consumption of crops grown on mine dumps which makes people vulnerable to these toxic elements.

References

- [1] Kabata-Pendias, H. (1999): *Trace Elements in Soil and Plants*, CRC Press, London, pg: 413.
- [2] Turner, P. (1994). The response of plants to heavy metals. In: Toxic metals in soil-plant system. Ross, M. (ed.), John Wiley & Sons Ltd., London, 153-182 pp.
- [3] Karczewskaa, K. (2002). Metale ciężkie w glebach zanieczyszczonych emisjami hut miedzi – formy i rozpuszczalność [Heavy metals in soils polluted by emissions from copper works – forms and solubility]. *Zesz. Nauk. AR Wroc.*, 432: 159 ss. (in Polish).
- [4] Wilson, B. and Pyatt, F. B. (2007). Heavy metal dispersion, persistence and bioaccumulation around an ancient copper mine situated Anglesey, UK. *Ecotoxicol Environ Saf* 66:224–231.
- [5] WHO (1992). Cadmium Environmental Health Criteria. World Health Organization, Geneva, p 134.
- [6] Steenland, K. and Boffetta, P. (2000). Lead and cancer in humans: where are we now? *Am. J. Ind Med* 38:295–299.
- [7] ATSDR (2008): *Minimal Risk Level*. Agency for Toxic Substances and Disease Registry. www.atsdr.cdc.gov. accessed on 01/12/2008.
- [8] Maleki, Z., Zaresvand (2008). Heavy metals in selected edible vegetables and estimation of their daily intake in Sanandaj, Iran. *Southeast Asian J. Trop Med Public Health* 39(2):335–340.
- [9] Lokeshappa, B, Shivpuri K, Tripath V, and Dikshit, A. K. (2012). Assessment of Toxic Metals in Agricultural Product. *Food Pub Health* 2(1):24–29.
- [10] Sresty, T. V. S., Rao KVM. (1999). Ultra structural alternation in response to zinc and nickel stress in the root cells of pea. *J Env Exp Bot* 41:3–13.
- [11] Lacatusu, R., Rauta, C., Carstea, S. and Ghelase, I. (1996). Soil - plant relationships in heavy metal polluted areas in Romania. *Appl Geochem* 11:105–107.
- [12] Page, A. L. (1991): *Methods of Soil Analysis*. 2nd edition. Am. Soc. Agron. & Soil Sci. Am. Madison, Wisconsin, USA.
- [13] Jarup, L. (2003). Hazards of Heavy Metal Contamination. *Br. Med. Bull.* 68:167–82.
- [14] Alloway, B. J. (1994): *Heavy Metals in Soils*. London, Blackie Academic and Professional. Pp: 368-400.